

R005-P08

ポスター 3 : 9/26 AM1/AM2 (9:00-12:30)

#津田 卓雄¹⁾, 安藤 芳晃¹⁾, 中川 広務²⁾, Ward William³⁾, 堤 雅基^{4,5)}, 穂積 裕太^{6,7)}, 細川 敬祐¹⁾, 村田 健史⁸⁾

(¹⁾ 電通大, (²⁾ 東北大, (³⁾ University of New Brunswick (UNB), (⁴⁾ 極地研, (⁵⁾ 総研大, (⁶⁾ National Aeronautics and Space Administration (NASA), (⁷⁾ Catholic University of America (CUA), (⁸⁾ 情報通信研究機構

Temperature retrieval in the middle atmosphere using Himawari-8/AHI limb-sounding data

#Takuo Tsuda¹⁾, Yoshiaki Ando¹⁾, Hiromu Nakagawa²⁾, William Ward³⁾, Masaki Tsutsumi^{4,5)}, Yuta Hozumi^{6,7)}, Keisuke Hosokawa¹⁾, Ken Murata⁸⁾

(¹⁾ University of Electro-Communications (UEC), (²⁾ Tohoku University, (³⁾ University of New Brunswick (UNB), (⁴⁾ National Institute of Polar Research (NIPR), (⁵⁾ Graduate University for Advanced Studies (SOKENDAI), (⁶⁾ National Aeronautics and Space Administration (NASA), (⁷⁾ Catholic University of America (CUA), (⁸⁾ National Institute of Information and Communications Technology (NICT)

Himawari-8 is the Japanese Geostationary Earth Orbit (GEO) meteorological satellite, that is equipped with Advanced Himawari Imager (AHI). Himawari-8/AHI provides full disk images every 10 min in 16 observation bands, including three visible bands: blue (0.47 μ m), green (0.51 μ m), and red (0.64 μ m). These full disk images are normally used as nadir observations mainly for meteorological purposes. On the other hand, the full disk images by Himawari-8/AHI can also provide limb-sounding data utilizing the edges of images with near-global coverage. As an example, there are a couple of reports on polar mesospheric cloud (PMC) observations by Himawari-8/AHI limb-sounding.

In the present work, we consider temperature retrieval in the middle atmosphere as a further application using Himawari-8/AHI limb-sounding data. In the limb-sounding, Rayleigh scattering of the sunlight can be observed, and thus we can obtain height profiles of line-of-sight (LOS) integrated Rayleigh scattering signals. By inversion methods, such as the Abel transforms, the onion peeling, etc., the LOS-integrated signals can be converted into local signals, which could be considered to be proportional to the local number densities of the atmospheric molecules. Then, applying the Rayleigh scattering temperature lidar technique, height profiles of temperature can be derived from height profiles of the local signals under an assumption of the hydrostatic equilibrium. There are a couple of previous works demonstrating this kind of temperature retrieval method using limb-sounding data from low-Earth-orbit (LEO) satellites. On the other hand, there is no previous example of GEO satellites, which have an advantage in providing continuous observations from a fixed point in space. In the presentation, we will show our initial data analysis for temperature retrieval in the middle atmosphere using Himawari-8/AHI limb-sounding data, and discuss the potential of the retrieved temperature data.